



Génétique cardiovasculaire: Concepts & applications

Jean-Jacques SCHOTT

Directeur de recherche INSERM

Equipe I « Génétique HUMAINE »

(jjschott@univ-nantes.fr)



L'unité de recherche de l'institut du thorax
Inserm UMR 1087 / CNRS UMR 6291
Nantes, France

MASTER 1 Thorax

Unité d'enseignement « Recherche en
Physiopathologies humaines »

17-11-2022

Approches génétique en 2022: objectifs

1- Concepts, outils & modèles: (Dynamique des génomes)

- Variants rares (→ ++ maladies mendéliennes)
 - Approches Liaison génétique – approches en famille (clonage positionnel)
 - Séquençage SANGER; séquençage d'exome (WES)
 - Séquençage de génomes complets (WGS) → accès à l'ensemble

du spectre de fréquences alléliques des REGIONS CODANTES & NON CODANTES; épigénétique)

- Variants fréquents
 - Etude d'association (GWAS); approches en population (Génotypage)
→ augmentation du risque

2- Applications en Génétique Cardiovasculaire




- Troubles de conduction cardiaque de l'enfance
- Dysfonction sinusale - cardiomyopathie
- Troubles du rythme ventriculaire: Le syndrome de Brugada

Strategy

Translational approach to investigate the genetic architecture of cardiovascular & developmental diseases

➤ **Identifying susceptibility genes and molecular mechanisms**

➤ **Largest collections worldwide**




New analytical tools:

Bioinformatics

Biostatistics

Epigenetics




Population

Common variants
(susceptibility genes)

Genotyping

Whole genome sequencing



Families


Rare variants
(strong effects)

Exome sequencing

Whole genome sequencing

NEW GENES

Functional studies



Disease gene identification allowed by a spectacular acceleration of knowledge of the architecture of the human genome:

Microsatellites

Genetic Maps



1990

2001

2010



SNPs



GWAS

WES

WGS

Rare diseases

(Rare diseases / Rare variants)

Common diseases


(Common diseases / common variants)



- new sequencing technologies
- new databases

- ... new disease models

Stratégies génétiques & dynamique des génomes



Novembre & Ramachandran.

Ann Rev Genomics Hum Genet **12**:245, 2011


- 1000 genome project (population)

- Bases de données : gnomAD, UK biobank (patients)

A global reference for human genetic variation

The 1000 Genomes Project Consortium*

- 2504 individuals from 26 different populations
- Broad spectrum of genetic variations:
 - 88 millions variants (MAF >1%)



bioRxiv is receiving many new papers on coronavirus SARS-CoV-2. A reminder: these are preliminary reports that have not been peer-reviewed and should not guide clinical practice/health-related behavior, or be reported in news media as established information.

New Results

[Comment on this paper](#)

High coverage whole genome sequencing of the expanded 1000 Genomes Project cohort including 602 trios

Marta Byrka-Bishop, Uday S. Evani, Xuefang Zhao, Anna O. Basile, Haley J. Abel, Allison A. Regier, André Corvelo, Wayne E. Clarke, Rajeeva Musunuri, Kshithija Nagulapalli, Susan Fairley, Alexi Runnels, Lara Winterkorn, Ernesto Lowy-Gallego, The Human Genome Structural Variation Consortium, Paul Flicek, Soren Germer, Harrison Brand, Ira M. Hall, Michael E. Talkowski, Giuseppe Narzisi, Michael C. Zody

doi: <https://doi.org/10.1101/2021.02.06.430068>

This article is a preprint and has not been certified by peer review [what does this mean?].

Variants rares

MAF (fréquence de l'allèle mineur) $< 0,1 \%$

Variations « récentes

Variants à effet fort

*Approche familiale –Maladies Mendéliennes
Souvent dans les gènes (exons)

Variants fréquents

MAF < 5 %

→ Variations « anciennes » maintenues durant l'évolution

→ Approche en population – Maladies fréquentes

« *Common variant – common disease* »


Variants à effet faible

Augmentation du risque de développer une maladie
d'environ 1- 5 %



Souvent dans les régions non codantes

Modulation de l'expression d'un gène

Clonage positionnel : « génétique des grandes familles » Variants rares




Clonage positionnel : Séquençage SANGER



Modes de transmission

Mode de transmission	Localisation du locus responsable du trait	Principales caractéristiques	Exemples de phénotypes
Autosomique dominant	Chromosome 1-22	-hommes et femmes sont atteints avec une fréquence équivalente -les enfants atteints ont un parent atteint	Huntington, syndrome de Marfan
Autosomique récessif	Chromosome 1-22	-hommes et femmes sont atteints avec une fréquence équivalente -les enfants atteints n'ont pas de parent atteint -peut « sauter » une ou plusieurs générations	Mucoviscidose, drépanocytose
Dominant lié à l'X	Chromosome X	-les femmes atteintes ont 50% d'enfants atteints -les hommes sont plus sévèrement atteints -toutes les filles de père atteints sont atteintes	Syndrome de Rett
Récessif lié à l'X	Chromosome X	-beaucoup plus d'hommes atteints que de femmes -les hommes atteints ont une mère saine (porteuse) -le trait n'est jamais transmis de père à fils -peut « sauter » une ou plusieurs générations	Hémophilie
Lié à l'Y	Chromosome Y	-tous les hommes atteints passent le trait à leur fils -les femmes ne sont pas atteintes	Stérilité
Mitochondrial	Pas chromosomique ADN mitochondrial	-tous les enfants d'une femme atteinte sont atteints -les hommes atteints ne transmettent pas le trait	Neuropathie optique de Leber

Effet des variants génétiques




- Effet de ces mutations variables selon la protéine
 - Haplo insuffisance (50 % de réduction d'activité)
 - Gain de fonction (ex. SCN5A /LQT3; fuite de courant sodique)
 - Effet dominant négatif (peptide poison; ex Canaux K⁺; structure $\alpha 4$)

Genetic variation detection improvement by the new technologies

Exome sequencing: capture all coding regions of the genome (1 - 2% soit 50 Mb)

nature



Analysis of 6,515 exomes reveals the recent origin of most human protein-coding variants

Wenqing Fu¹, Timothy D. O'Connor¹, Goo Jun², Hyun Min Kang², Goncalo Abecasis², Suzanne M. Mark J. Rieder¹, David Altshuler⁴, Jay Shendure¹, Deborah A. Nickerson¹, Michael J. Bamshad^{1,5}, N. & Joshua M. Akey¹






Evolution and Functional Impact of Rare Coding Variation from Deep Sequencing of Human Exomes

Jacob A. Tennessen *et al.*

Science 337, 64 (2012);

DOI: 10.1126/science.1219240



Séquençage d'Exome (NGS)



Reference	..TTTAGTAATAACAT--GTGCATCTTTAATC..
Read_01	..TTTAGGAATAACATGCGTGCA
Read_02	..TTTAGGAATAACATGCGTGCATCT
Read_03	..TTTAGGAATAACATGCGTCCATCTT
Read_04	..TTTAGGAATAACATGCGTGCATCTTTAAT
Read_05	..TTTAGGAATAACATGCGTGCATCTTTAATC..
Read_06	..TTTAGGAATAACATGCGTGCATCTTTAATC..
Read_07	..TTTAGGAATAACATGCGTGCATCTTTAATC..
Read_08	AATAACATGCGTGCATCTTTAATC..
Read_09	TAACATGCGTGCATCTGTAATC..

Séquengage « Sanger »


Sample → FASTQ




```
@SRR038845.3 HWI-EAS038:6:1:0:1938 length=36
CAACGAGTTCACACCTTGGCCGACAGGCCCGGGTAA
+SRR038845.3 HWI-EAS038:6:1:0:1938 length=36
BA@7>B=>:>>7@7@>>9=BAA?;>52;>:9=8.=A
@SRR038845.41 HWI-EAS038:6:1:0:1474 length=36
CCAATGATTTTTTTTCCGTGTTTCAGAATACGGTTAA
+SRR038845.41 HWI-EAS038:6:1:0:1474 length=36
BCCBA@BB@BBBBBAB@B9B@=BABA@A:@693:@B=
@SRR038845.53 HWI-EAS038:6:1:1:360 length=36
GTTCAAAAAGAACTAAATTGTGTCAATAGAAAACCTC
+SRR038845.53 HWI-EAS038:6:1:1:360 length=36
BBCBBBBBBB@@BAB?BBBBBCBC>BBBAA8>BBBAA@
```

FASTQ → BAM


```
@SRR038845.3 HWI-EAS038:6:1:0:1938 length=36
CAACGAGTTTACACCTTGGCCGACAGGCCCGGGTAA
+SRR038845.3 HWI-EAS038:6:1:0:1938 length=36
BA@7>B=>:>7@7@>9=BAA?;>52;>9=8.=A
@SRR038845.41 HWI-EAS038:6:1:0:1474 length=36
CCAATGATTTTTTCCGTGTTTCAGAATACGGTTAA
+SRR038845.41 HWI-EAS038:6:1:0:1474 length=36
BCCBA@BB@BBBBBAB@B9B@=BABA@A:@693:@B=
@SRR038845.53 HWI-EAS038:6:1:1:360 length=36
GTTCAAAAAGAACTAAATTGTGTCAATAGAAAACCTC
+SRR038845.53 HWI-EAS038:6:1:1:360 length=36
BBCBBBBBB@B@BAB?BBBBBCBC>BBBAA8>BBBAA@
```



BAM




BAM → VCF




CHROM	POS	REF	S1	S2
1	1	A	A/A	A/A
12	234	T	T/A	T/T
20	34	G	G/A	G/G
X	344	C	G/A	C/C

VCF+Annotations



Variant Filtering



gnomAD browser beta | genome Aggregation Database

Search for a gene or variant or region

Example - Gene: [PCSK9](#), Variant: [1-55516888-G-GA](#)

About gnomAD

The [Genome Aggregation Database](#) (gnomAD) is a resource developed by an international coalition of investigators, with the goal of aggregating and harmonizing both exome and genome sequencing data from a wide variety of large-scale sequencing projects, and making summary data available for the wider scientific community.

The data set provided on this website spans 123,136 exome sequences and 15,496 whole-genome sequences from unrelated individuals sequenced as part of various disease-specific and population genetic studies. The gnomAD Principal Investigators and groups that have contributed data to the current release are listed [here](#).

All data here are released for the benefit of the wider biomedical community, without restriction on use - see the terms of use [here](#).

Recent News

February 27, 2017

[Official gnomAD release \(version 2.0\)](#) with browser updates and data available for [download](#).

October 19, 2016

Public release of gnomAD Browser (beta) at ASHG!

Identification de variants rares causaux grâce aux nouvelles technologies de séquençage



1. Variants fonctionnels

Variants non-synonymes et d'épissage

2. Variants rares

Bases de données externes et internes

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
10	5138672	C	G

~ 85.000 variants

22	16449042	T	A
5	56177051	C	G
5	75913610	C	T
6	10894103	C	T
9	102068277	A	C
9	112015730	T	C
1	897829	C	G
1	8415627	T	C
1	17722114	G	A
1	86173964	C	T
1	92765756	A	G
1	113238173	C	T
1	120458982	C	G
1	143378919	T	C
1	146639425	G	A
1	159002410	A	G
1	159021815	T	G
1	177250522	G	A
1	179012995	CAGCG	C
1	242022009	G	A
1	247463963	C	T
1	247492516	C	T
10	17083054	T	C
10	28233186	G	A
10	83635880	A	G
10	83992548	G	T
10	99377056	G	T
10	102265186	C	T
10	102778027	TCTC	T
10	116059031	G	C
10	121347832	C	A
10	127429028	A	G
11	1298431	G	A
11	1463811	G	A
11	1904753	A	G
11	3681258	G	C
11	6636687	G	A
11	27363105	A	C
11	34527245	C	T
11	46401474	G	A
11	55111661	TAA	T
11	55433325	C	T
11	59712732	G	A
11	62287110	C	T
11	63066440	T	G
11	64428387	C	T
11	67265077	C	T
11	73744718	C	A

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
10	5138672	C	G

~ 15 000 variants


22	16449042	T	A
5	56177051	C	G
5	75913610	C	T
6	10894103	C	T
9	102068277	A	C
9	112015730	T	C
1	897829	C	G
1	8415627	T	C
1	17722114	G	A
1	86173964	C	T
1	92765756	A	G
1	113238173	C	T
1	120458982	C	G
1	143378919	T	C
1	146639425	G	A
1	159002410	A	G
1	159021815	T	G
1	177250522	G	A
1	179012995	CAGCG	C
1	242022009	G	A
1	247463963	C	T
1	247492516	C	T

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
10	5138672	C	G

~ 200 variants

18	31324584	G	A
1	247492516	C	T

Identification de variants rares causaux grâce aux nouvelles technologies de séquençage



1. Variants fonctionnels

Variants non-synonymes et d'épissage

2. Variants rares

Bases de données externes et internes

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
10	5138672	C	G

~ 85.000 variants

22	16449042	T	A
5	56177051	C	G
5	75913610	C	T
6	10894103	C	T
9	102068277	A	C
9	112015730	T	C
1	897829	C	G
1	8415627	T	C
1	17722114	G	A
1	86173964	C	T
1	92765756	A	G
1	113238173	C	T
1	120458982	C	G
1	143378919	T	C
1	146639425	G	A
1	159002410	A	G
1	159021815	T	G
1	177250522	G	A
1	179012995	CAGCG	C
1	242022009	G	A
1	247463963	C	T
1	247492516	C	T
10	17083054	T	C
10	28233186	G	A
10	83635880	A	G
10	83992548	G	T
10	99377056	G	T
10	102265186	C	T
10	102778027	TCTC	T
10	116059031	G	C
10	121347832	C	A
10	127429028	A	G
11	1298431	G	A
11	1463811	G	A
11	1904753	A	G
11	3681258	G	C
11	6636687	G	A
11	27363105	A	C
11	34527245	C	T
11	46401474	G	A
11	55111661	TAA	T
11	55433325	C	T
11	59712732	G	A
11	62287110	C	T
11	63066440	T	G
11	64428387	C	T
11	67265077	C	T
11	73744718	C	A

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
10	5138672	C	G


~ 15 000 variants

22	16449042	T	A
5	56177051	C	G
5	75913610	C	T
6	10894103	C	T
9	102068277	A	C
9	112015730	T	C
1	897829	C	G
1	8415627	T	C
1	17722114	G	A
1	86173964	C	T
1	92765756	A	G
1	113238173	C	T
1	120458982	C	G
1	143378919	T	C
1	146639425	G	A
1	159002410	A	G
1	159021815	T	G
1	177250522	G	A
1	179012995	CAGCG	C
1	242022009	G	A
1	247463963	C	T
1	247492516	C	T

CRHOM	POSITION	REF	ALT
5	112175526	G	C
5	125820120	G	C
18	31324584	G	A
1	247492516	C	T

~ 200 variants

➤ Partagés par les individus atteints (cas familiaux)



➤ Liaison génétique

Classification du niveau de pathogénicité

1- Méthodes « *in silico* » de classification de variants génétiques

- ✓ Variants localisés dans une séquence codante

SIFT, Polyphen-2, Crescendo, Bongo, SDM, MutPred, PoPMusic, PHD-SNP, Mu-Pro...

- ✓ Variants localisés dans une séquence non-codante

Eponine, McPromoter, PromoterScan, Match/Transfac, TFSEARCH, ESEFinder, RESCUE-ESE...
+ données ENCODE (Septembre 2012)


2- Méthodes « *in vitro* »

- tests fonctionnels (bas-débit → Patch clamp (canaux ioniques); souris transgéniques (CRISPR/Cas9)
- évolution technologique:
 - patch haut-débit
 - modèles cellulaires & animaux : cellules souches pluripotentes induites (IPS); poisson zèbre)







Classification ACMG (American College of Medical Genetics)

- Classe 1 -5: Score de niveau de confiance de pathogénicité:
pathogenic, likely pathogenic, uncertain significance, likely benign, and benign
- VSI: Variants de signification inconnue


Identification par séquençage d'exomes / génomes de variants rares



Genetics of primary electrical cardiac disorders



Primary electrical cardiac disorders




Sick sinus syndrome


Atrial arrhythmias
Atrial Fibrillation

Conduction defects
Atrioventricular block


Ventricular arrhythmias
Long QT syndrome
Brugada Syndrome
Early Repolarisation



Probst et al., 2009



Primary electrical cardiac disorders




Sick sinus syndrome

Atrial arrhythmias
Atrial Fibrillation


Conduction defects
Atrioventricular block

Ventricular arrhythmias
Long QT syndrome
Brugada Syndrome
Early Repolarisation




■ BrS
● Conduction disease
□ Unaffected

Probst et al.,
2009




..... Rare to common cardiac diseases

Rare and Common variants associated with cardiac arrhythmia present in coding and non-coding region



Genome-wide association analyses identify new Brugada syndrome risk loci and highlight a new mechanism of sodium channel regulation in disease susceptibility

Rare and Common variants associated with cardiac arrhythmia present in coding and non-coding region



Genome-wide association analyses identify new Brugada syndrome risk loci and highlight a new mechanism of sodium channel regulation in disease susceptibility

CTCF-mediated chromatin looping disruption causes PITX2-related novel cardiac syndrome

Identification of a Deletion Upstream SCN5A associated with Brugada syndrome

De novo mutations in congenital cardiac conduction defects

Common non-coding variant

Rare non-coding variant

Rare coding variant

2023 : Cardiac arrhythmias genetics (> 60 genes)




Long QT Syndrome (15)

KCNQ1
KCNH2
SCN5A
ANK2
KCNE1
KCNE2

KCNJ2
KCNJ5
CACNA1C
CAV3
SCN4B
AKAP9
SNTA1

CALM1




Brugada Syndrome (25)

SCN5A
GPDI1L
CACNA1C
CACNB2
SCN1B

SCN3B
KCND3
KCNH2
HCN4
KCNJ8
SLMAP
CACNA2D1
RANGRF
SCN2B
ABCC9 ...


TRPM4

KCNE5
KCNE3
KCNAB2
RRAD



Short QT

KCNQ1
KCNH2
KCNJ2
CACNA1C
CACN2B
CACNA2D1




Atrial Fibrillation

KCNQ1
ANK2
ABCC9
GJA5
KCNE1
KCNE2
KCNJ2

SCN5A
SCN3B
SCN1B
...
SCN2B


KCNE5
KCNA5



Early repolarization Syndrome

KCNJ8
SCN5A
CACNA1C

CACNB2
CACNA2D1
ABCC9



CPVT


RYR2
CASQ2
TRDN
CALM1

- Familial rare (linkage/candidate gene)
- Overlap syndrome
- Variable penetrance and expressivity



Progressive conduction (Lenègre)

SCN5A
SCN1B
TRPM4
GJA5
GJA1



ACMG Recommendations for Reporting of Incidental Findings in Clinical Exome and Genome Sequencing

The American College of Medical Genetics and Genomics has published recommendations for reporting incidental findings in the exons of certain genes.


Adenomatous polyposis coli (MIM 175100)	APC (MIM 611731)
Aortic aneurysm, familial thoracic 4 (MIM 132900)	MYH11 (MIM 160745)
Aortic aneurysm, familial thoracic 6 (MIM 611788)	ACTA2 (MIM 102620)
Arrhythmogenic right ventricular cardiomyopathy, type 5 (MIM 604400)	TMEM43 (MIM 612048)
Arrhythmogenic right ventricular cardiomyopathy, type 8 (MIM 607450)	DSP (MIM 125647)
Arrhythmogenic right ventricular cardiomyopathy, type 9 (MIM 609040)	PKP2 (MIM 602861)
Arrhythmogenic right ventricular cardiomyopathy, type 10 (MIM 610193)	DSG2 (MIM 125671)
Arrhythmogenic right ventricular cardiomyopathy, type 11 (MIM 610476)	DSC2 (MIM 125645)
Breast-ovarian cancer, familial 1 (MIM 604370)	BRCA1 (MIM 113705)
Breast-ovarian cancer, familial 2 (MIM 612555)	BRCA2 (MIM 600185)
Brugada syndrome 1 (MIM 601144)	SCN5A (MIM 600163)
Catecholaminergic polymorphic ventricular tachycardia (MIM 604772)	RYR2 (MIM 180902)
Dilated cardiomyopathy 1A (MIM 115200)	LMNA (MIM 150330)
Dilated cardiomyopathy 1A (MIM 115200)	MYBPC3 (MIM 600958)
Ehlers-Danlos syndrome, type 4 (MIM 130050)	COL3A1 (MIM 120180)
Fabry's disease (MIM 301500)	GLA (MIM 300644)
Familial hypercholesterolemia (MIM 143890)	APOB (MIM 107730)
	LDLR (MIM 606945)
Familial hypertrophic cardiomyopathy 1 (MIM 192600)	MYH7 (MIM 160760)
Familial hypertrophic cardiomyopathy 3 (MIM 115196)	TPM1 (MIM 191010)
Familial hypertrophic cardiomyopathy 4 (MIM 115197)	MYBPC3 (MIM 600958)
Familial hypertrophic cardiomyopathy 6 (MIM 600858)	PRKAG2 (MIM 602743)
Familial hypertrophic cardiomyopathy 7 (MIM 613690)	TNNI3 (MIM 191044)
Familial hypertrophic cardiomyopathy 8 (MIM 608751)	MYL3 (MIM 160790)
Familial hypertrophic cardiomyopathy 10 (MIM 608758)	MYL2 (MIM 160781)
Familial hypertrophic cardiomyopathy 11 (MIM 612098)	ACTC1 (MIM 102540)
Familial medullary thyroid carcinoma (MIM 155240)	RET (MIM 164761)
Hypercholesterolemia, autosomal dominant, 3 (MIM 603776)	PCSK9 (MIM 607786)
Juvenile polyposis syndrome, (MIM 174900)	BMPR1A (MIM 601299)
Juvenile polyposis syndrome, (MIM 174900)	SMAD4 (MIM 600993)
Left ventricular noncompaction 6 (MIM 601494)	TNNT2 (MIM 191045)
Li-Fraumeni syndrome 1 (MIM 151623)	TP53 (MIM 191170)
Loeys-Dietz syndrome type 1A (MIM 609192)	TGFBR1 (MIM 190181)
Loeys-Dietz syndrome type 1B (MIM 610168)	TGFBR2 (MIM 190182)
Loeys-Dietz syndrome type 2A (MIM 608967)	TGFBR1 (MIM 190181)
Loeys-Dietz syndrome type 2B (MIM 610380)	TGFBR2 (MIM 190182)
Loeys-Dietz syndrome type 3 (MIM 613795)	SMAD3 (MIM 603109)
Long QT syndrome 1 (MIM 192500)	KCNQ1 (MIM 607542)
Long QT syndrome 2 (MIM 613688)	KCNH2 (MIM 152427)
Long QT syndrome 3 (MIM 603830)	SCN5A (MIM 600163)

➤ 78 actionable genes


■ 19 cardiac diseases

- 15 cardiomyopathies (ARVC, DCM, HCM)
- 4 arrhythmias

- Brugada Syndrome




- Long QT Syndrome



Cardiac Conduction Defects



- Historically isolated Progressive Cardiac Conduction Defects, the most common form of CCD, (Lenègre & Lev disease) was considered as a structural and degenerative disease mostly due to **aging**, fibrosis in the conduction system



◆ Strong genetic background / genetic heterogeneity:

- Isolated PCCD: Channelopathy, *SCN5A*, *SCN1B*, *TRPM4*, *GJA5*
- PCCD & Cardiomyopathies: *NKX2.5*, *TBX5*, *PRKAG2*, *LMNA*
- Non-immune isolated Congenital AV block is a rare condition
 - Prevalence : 1/15 000
 - Immunological compound in 80 % congenital AVB
 - Mortality : 30 %




◆ Suspected Strong genetic background

Search for new Congenital & early childhood AVB index genes: Strategy

- Patient recruitment:
 - 141 Index cases
 - Diagnosed under age 15 (13 centers in France)

➤ cAVB – trios



- 15 trios
 - No familial history of AV block
 - Unaffected parents

➤ Exome sequencing



European Heart Journal Advance Access published September 14, 2011

European Heart Journal
doi:10.1093/eurheartj/ehs347

CLINICAL RESEARCH

Characteristics and long-term outcome of non-immune isolated atrioventricular block diagnosed *in utero* or early childhood: a multicentre study

Alban-Elouen Baruteau^{1,2,3,4,5,6*}, Swanny Fouchard¹, Albin Behaghel^{2,3,4,7}, Philippe Mabo^{2,3,4,7}, Elisabeth Villain^{6,8}, Jean-Benoit Thambo^{6,9}, François Marçon^{6,10}, Véronique Gournay^{6,11}, Francis Rouault^{6,12}, Alain Chantepie^{6,13}, Sophie Guillaumont^{6,14}, François Godart^{6,15}, Caroline Bonnet^{6,16}, Alain Fraisse^{6,17}, Jean-Marc Schleich^{2,3,4,6,7}, Jean-René Lussan^{6,18}, Yves Dulac^{6,19}, Christophe Leclercq^{2,3,4,7}, Jean-Claude Daubert^{2,3,4,7}, Jean-Jacques Schott¹, Hervé Le Marec^{1,20}, and Vincent Probst^{1,20}

 Agilent Technologies



Exome sequencing of 15 trios

➤ 19 *de novo* variants

Trio	Gène	Nucléotide	Acide Aminé	Conséquence
0	<i>NSUN6</i>	c.956 C>T	p.R319Q	Faux-sens
1	<i>GCN1L1</i>	G>T		Site d'épissage (intronique)
2	<i>SHBG</i>	c.74G>A	p.R25H	Faux-sens
2	<i>CNTROB</i>	c.137G>A	p.R46Q	Faux-sens
3	<i>USH2A</i>	c.82T>A	p.I28L	Faux-sens
3	<i>SOS2</i>	c.58G>C	p.R20G	Faux-sens
4	<i>GPR107</i>	c.1396C>A	p.L466I	Faux-sens
6	<i>PTGER3</i>	C>A		Site d'épissage (intronique)
7	<i>TTC21B</i>	c.3641 T>C	p.D1214G	Faux-sens
7	<i>C7orf61</i>	c.209 A>G	p.L70S	Faux-sens
8	<i>TEX29</i>	G>C		Site d'épissage (5'-UTR)
9	<i>LPHN2</i>	c.296 A>G	p.N99S	Faux-sens
9	<i>RHOBTB1</i>	c.1259 G>A	p.T420M	Faux-sens
10	<i>RYR2</i>	c.10361 G>A	p.R3454H	Faux-sens
11	<i>ZNF683</i>	c.1219_1226 delCTGCACTG	p.Q407CfsX461	Frameshift & protéine tronquée
11	<i>ZNF22</i>	c.85 C>T	p.Q29X	Gain d'un codon-stop
11	<i>GJC1</i>	c.224 C>T	p.R75H	Faux-sens
11	<i>ATAD2B</i>	c.4212 C>G	p.L1404F	Faux-sens
14	<i>DOPEY2</i>	c.6697 C>T	p.R2233C	Faux-sens



Variant selection




	Family A (trio11)	Family B
	French case	Japanese case
Target genes for exon capture	All exons (Trio-exome)	457 genes
Coding variations (*)	13,035	1,040
<i>De novo</i> variants	7	NA
MAF<0.1%	4	8
Sanger validation	4	8
Co-segregation	NA	3
Candidate variations	4	3
Common mutation	<i>GJC1</i> -c.G224A, (Cx45-p.R75H)	

*: nonsynonymous, stop-gain, stop-loss, splice site replacement, or indel frameshift


NA: not applicable

¶; found in the proband (II:1) but not in his parents

Novel *GJC1* mutation (Cx45-p.R75H)




		64	E1	75	TM2	87
Cx45 Human		CYDAFAPLSHV	R	FWVFQIILVATP		
Chimp						
Macaca						
Mouse						
Rat						
Dog						
Opossum						
Cx43 Human		...KSF.I...	.	...L...F.SV.		
Cx40 Human		...QAF.I..I	I	.Y..L...F.S..		
Cx26 Human		...HYF.I..I	.	L.AL.L.F.S..		
Cx32 Human		...Q.F.I...	.	L.SL.L...S..		
Cx46 Human		...RAF.I..I	.	...AL...F.S..		




- Connexin family of proteins consists of more than **21 members** varying in their biophysical properties
- **CX40, 43, 45** heart specific


Family A




II:1
(2 yo)



(4 yo)



(5 yo)




(9 yo)




3rd degree AV block & progressive loss of P wave


Family B



II:2
(9 yo)




III:1
(8 yo)




(12 yo)




(14 yo)



III:2
(14 yo)




I:1
(37 yo)




In vitro studies

1- Normal hemichannel assembly and plaque formation


(Co-IP /Immunofluorescent)




(Immunofluorescent)



2- Severe reduction of permeation properties (Luciferase dye transfer)



3- Macroscopic conductance (Gj) suppressed in hetero & homomeric channels




➤ Dominant-negative

In vivo studies



Electrophysiological properties of conditional *Gjc1* knockout mice



1- Various atrial
arrhythmias
(after tamoxifen
administration)




➤ Pathophysiology of Cx45-p.R75H in developmental defects?

Seki et al. J Am Coll Cardiol. 2017


Whole Genome Sequencing in unresolved mendelian arrhythmias



A large family associating long QT syndrome and sinus dysfunction linked to the ANK2 gene (LQT4)




➤ Loss of function mutation in *ANK2* (*Ankyrin*)



Mohler P; Schott JJ et al. Nature. 2003

- 24 affected cases
- 2 sudden death (ventricular arrhythmias)




- Long QT syndrome (LQT4) T/U waves
- Sinus node dysfunction (SND)
- Atrial fibrillation (AF)

➤ Additional long QT / SND *ANK2*^{-/-} family but with genetic linkage at 4q25 locus

➤ 4q25 linkage

4




Le Scouarnec et al. Proc Natl Acad Sci. 2008

Schott et al. Am J Hum Genet. 1995

Identification of a large family presenting with a complex cardiac syndrome

17 members present with a complex cardiac syndrome following an autosomal dominant model

Family 1



Electrical cardiac disorders:


Sinus node dysfunction
Long QT
Atrial fibrillation

Cardiac developmental defect:

Atrial septal defect
LV non-compaction

Identification of a large family presenting with a complex cardiac syndrome associated with a 15 kb deletion in a gene desert area

Family 1



Electrical cardiac disorders:

Sinus node dysfunction
Long QT
Atrial fibrillation

Cardiac developmental defect:

Atrial septal defect
LV non-compaction

Linkage analysis

4q25 region associated with
Ankyrin B syndrome
PJ.Molher et al 2002

Sanger sequencing


No mutation in *ANK2* gene

Exome sequencing

No mutation in genes located in
the 4q25 region

Whole genome sequencing


Identification of a 15kb
deletion in a gene desert area



Reduced sequencing depth by 50%

Identification of a large family presenting with a complex cardiac syndrome associated with a 15 kb deletion in a gene desert area

Family 1



Electrical cardiac disorders:

Sinus node dysfunction
Long QT
Atrial fibrillation

Cardiac developmental defect:

Atrial septal defect
LV non-compaction

PITX2

- Cardiac transcription factor essential for the left-right asymmetry of the heart
- SA development
- Risk variants in *PITX2* associated with atrial fibrillation (GWAS)

Ank2

- Localization and membrane stabilization of ion channels
- *ANK2* loss-of-function variants -> LQT4 syndrome (LQT-SND)



Human chr4

1 Mb

hg19

refSeq genes

ENPEP

PITX2

Family1 15 kb

C4orf32

AP1AR

TIFA

ALPK1

NEUROG2

C4orf21


LARP7

ANK2

CAMK2D

Relevance of this non-coding region in the molecular mechanisms leading to this new syndrome ?


Six additional families (4 French and 2 Japanese) presenting with a similar phenotype showing overlapping deletions



Human chr4

refSeq genes

ENPEP
PITX2



C4orf32
AP1AR
TIFA
ALPK1
NEUROG2
C4orf21
LARP7

ANK2
CAMK2D

Crucial role of this non coding deleted region

Molecular mechanism


In vitro



iPS isogenic model

iPSC differentiation to cardiomyocytes & genome editing


iPS-CM



Functional studies


- Cell morphology
- Cell beat recording
- Cardiomyocytes contractility
- Action potential recording

Chromatin conformation




MAX-PLANCK-GESELLSCHAFT


In vivo



Human chr4 1 Mb hg19
refSeq genes
ENPEP
PITX2




Mouse Chr3 mm10
refSeq genes
Pitx2
Enpep



Epigenetics: Understanding the non-coding regions of the human genome

Every cell-type in an organism contains the same genetic information but has extremely different shapes and functions.




How cells read the genes

<https://www.slideshare.net/hamivia/definition-of-epigenetics>

Definition of Epigenetics

Every cell-type in an organism contains the same genetic information but has extremely different shapes and functions.



How cells read the genes

<https://www.slideshare.net/hamivia/definition-of-epigenetics>


DNA methylation

Chromatin structure and modification

Non-coding RNA (ncRNA)


Definition of Epigenetics

Every cell-type in an organism contains the same genetic information but has extremely different shapes and functions.




How cells read the genes

DNA methylation
Chromatin structure and modification
Non-coding RNA (ncRNA)






Definition of Epigenetics

Every cell-type in an organism contains the same genetic information but has extremely different shapes and functions.





How cells read the genes




Tissue-specific

Species-specific



Non-coding region characterization and annotation




How cells read the genes

<https://www.slideshare.net/hamivia/definition-of-epigenetics>

ATAC-seq approach: assay for Transposase-Accessible Chromatin





ChIP-seq: chromatin immunoprecipitation –




ChIP-Seq mark	Functional association
H3K4me3	promoters
H3K4me1	enhancers
H3K27ac	Active promoters and enhancers
H3K27me3	Inactive chromatin






Functional annotation of non-coding region in the 4q25 region










Functional annotation of non-coding region in the 4q25 region




Hypothesis: Deletion of the CTCF binding site modifies the TAD boundary




Deletion of CTCF
binding site



New regulatory region interacting with *PITX2*



Mouse model to characterize the impact of the deletion on the phenotype



Human chr4

1 Mb hg19

RefSeq genes

ENPEP
PITX2

Family 1 15 kb
Family 2 32 kb
Family 3 47 kb
Family 4 92 kb
Family 5 146 kb
Family 6 150 kb
Family 7 330 kb

C4orf32

NEUROG2

ANK2

CAMK2D

AP1AR

C4orf21

TIFA

LARP7

ALPK1

Human-mouse alignable

Mouse Chr3

mm10 1 Mb

RefSeq genes

Pitx2
Enpep

Tifa

Zgrf1

Ank2

Camk2d


Ap1ar

Larp7





Fam241a

Neurog2

Alpk1



Mouse model to characterize the impact of the deletion on the phenotype



No phenotype observed in mice carrying deletion A and C



In vivo electrical cardiac activity in mice carrying the 15kb deletion

Mouse Chr3 mm10 1 Mb

refSeq genes

Pitx2
Enpep


Tifa *Zgrf1* *Ank2*
Ap1ar *Larp7*
Fam241a *Neurog2* *Camk2d*
Alpk1



CRISPR mice show slower heart rates, slower sinus node recovery times, atrial arrhythmias



Bradycardia, SND and atrial arrhythmia recapitulates electrical phenotype identified in patients

Cardiac morphological characterization in mice carrying the 15kb deletion



CRISPR mice present ASD

Transcriptomic analysis



Upregulation of *Pitx2* in the right atria



Downregulation of *Pitx2* & *Ank2* in the ventricles

Manuscript in preparation



Overexpression of *PITX2* during differentiation of sinoatrial nodal cells


CTCF- KO (19 bp)




Conclusions & Perspectives

- **WGS & multi-omics are powerful approaches to identify new regulatory elements and target genes in unsolved mendelian diseases**
 - New regulatory mechanism impacting *PITX2* expression
 - New role of *PITX2* in disease: new cardiac syndrome associating atrial, ventricular arrhythmias and congenital heart defects (ASD or LVNC)
- **Implications for clinical molecular diagnosis**
 - Search for 4q25-CTCF deletion should firstly be performed in patients presenting a combination of electrical (LQT/SND) and structural heart defects phenotypes
 - Implication of 4q25-CTCF deletions in isolated congenital heart defects is ongoing

Fibrillation ventriculaire idiopathique



Mort subite cardiaque: population à risque



Cardiac arrhythmia

➤ **Sudden Cardiac Death is a major health burden in industrialized countries**


➤ *Ventricular Fibrillation is the most common mechanism*

Myocardial infarction

Cardiomyopathy

~80%

~10%



Acute plaque destabilization



Hypertrophic Cardiomyopathy




Dilated Cardiomyopathy



Primary electrical disorders

~10%



- ⇒ **Brugada syndrome**
- ⇒ **Early repolarization syndrome**
- ⇒ **Long QT Syndrome**
- ⇒ **Catecholaminergic Polymorphic Ventricular Tachycardia**
- ⇒ **Arrhythmogenic right ventricular cardiomyopathy**
- ⇒ **Sinus node dysfunction**
- ⇒ **Cardiac conduction defects**



These electrical disorders provide a homogenous groups of patients with high risk for SCD, which can be considered as a 'sensitized model' for VF

The Brugada syndrome

- ◆ Rare disease (1-5/10 000), affecting mainly men ($\approx 80\%$)
- ◆ Mean age of diagnosis around age 40
- ◆ Ventricular tachycardia, ventricular fibrillation and sudden cardiac death
- ◆ Baseline condition or after drug challenge
Typical pattern of ST segment elevation, on the right precordial leads (V1-V3) of the surface ECG
- ◆ No cardiac structural defects

◆ Brugada syndrome is a genetic condition:

- Autosomal dominant disease
- **SCN5A** loss of function mutations in **$\sim 20\%$ cases**
 - Decreased upstroke velocity of the cardiac action potential
 - Reduced penetrance and variable expressivity
- 22 additional genes (familial and candidate gene approaches)
- Over all contribution to disease prevalence is still unclear
- **Burden testing:** only rare **SCN5A** variants are significantly associated to BrS



➤ **SCN5A** remains a poor marker for SCD risk stratification (BrS, a polygenic disease?)

ORIGINAL ARTICLE

Testing the burden of rare variation in arrhythmia-susceptibility genes provides new insights into molecular diagnosis for Brugada syndrome

Gene	BrS cases (n = 167)	Internal controls (n = 167)	P-value 1	UK10K controls (n = 881)	P-value 2
BrS-susceptibility genes					
SCN5A	20.4% (34)	2.4% (4)	1.4×10^{-7a}	2.4% (21)	1.7×10^{-15a}
SCN10A	6% (10)	2.4% (4)	0.170	3.5% (31)	0.131
CACNA1C	3% (5)	6.6% (11)	0.199	2% (18)	0.395
PKP2	3% (5)	2.4% (4)	1	1.7% (15)	0.348
CACNB2	1% (3)	1.2% (2)	1	0.9% (8)	0.396
KCNH2	1.2% (2)	3.6% (6)	0.283	1.6% (14)	1
TRPM4	1.2% (2)	3% (5)	0.448	1.9% (17)	0.754
KCND3	0.6% (1)	1.2% (2)	1	1.6% (14)	0.488
CACNA2D1	0.6% (1)	0.6% (1)	1	3.3% (29)	0.072
HEY2	0.6% (1)	0.6% (1)	1	0.1% (1)	0.293
SCN2B	0.6% (1)	0.6% (1)	1	0.5% (4)	0.581
SCN3B	0.6% (1)	0.6% (1)	1	0.5% (4)	0.581
ABCC9	—	3% (5)	0.061	1.1% (10)	0.379
SCN1B	—	1.8% (3)	0.248	0.3% (3)	1
RANGRF	—	0.6% (1)	1	0.2% (2)	1
FGF12	—	—	—	0.7% (6)	0.597

 Nous ne pouvons pas afficher l'image.


Cases

Controls

Significant enrichment in rare coding variation for *SCN5A* only

Variable penetrance and expressivity in mendelian diseases

Brugada Syndrome families with non-mendelian segregation



Presence of genetic modulators?


➤ Role of frequent variants modulating BrS expressivity

Common polymorphisms modulate Brugada syndrome


L'institut du thorax
in the heart of an international network on Brugada syndrome





312 patients
1115 controls




*** National Referral Centre for Inherited Cardiac Arrhythmias**




Cumulative effect of risk alleles at the three loci



■ Ctrl
□ BrS cases



International Brugada Syndrome Genetics Consortium




**~2800 BrS probands,
European descent**

Brugada syndrome Genome-wide association meta-analysis



GWAS: identification of 12 loci associated with Brugada syndrome

21 independent association signals (10 new loci)




New molecular mechanisms

➤ BrS heritability (h^2):

- ◆ substantial portion of susceptibility to BrS attributable to common genetic variation
- ◆ 0.17 (LDSC) to 0.34 (GREML)

A predominant role of *SCN5A/SCN10A* locus risk alleles



➤ *SCN5A/SCN10A* locus: 8 independent loci


SCN5A/SCN10A locus: 8 independant loci

Locus	Lead SNP	Genomic position (hg19)	Risk allele	Other allele	Risk allele frequency in cases	Risk allele frequency in controls	OR [95% CI]	P value	Nearest gene
1	rs7638909*	3:38594973	G	T	0.32	0.24	1.28 [1.17 - 1.40]	2.79E-08	SCN5A
	rs62241190*	3:38607468	G	A	0.06	0.03	1.96 [1.63 - 2.32]	8.56E-14	SCN5A
	rs7374540*	3:38634142	C	A	0.51	0.39	1.72 [1.61 - 1.81]	3.56E-57	SCN5A
	rs7433206*	3:38657708	A	T	0.45	0.42	1.48 [1.37 - 1.60]	9.52E-24	SCN5A
	rs34760424*	3:38683018	G	T	0.98	0.94	2.32 [1.96 - 2.70]	3.03E-23	SCN5A
	rs41310232*	3:38689242	A	G	0.16	0.09	1.56 [1.40 - 1.74]	1.19E-15	SCN5A
	rs6782237*	3:38696553	C	G	0.78	0.68	1.74 [1.61 - 1.87]	1.05E-47	SCN5A
	rs6801957	3:38767315	T	C	0.65	0.42	2.49 [2.34 - 2.65]	1.30E-180	SCN10A

➤ All non-coding risk alleles: role in gene regulation


- ◆ post-GWAS approaches: translate genetic findings into molecular mechanisms
- ◆ characterize and annotate all cardiac regulatory regions

Risk haplotypes co-localize at multiple SCN10A / SCN10A intronic enhancers



- Epigenetic marks:
- Histone acetylation, methylation
- TF bindings sites
- 3D chromatin conformation


GWAS: identification of 8 loci associated with transcription factors




➤ Implication of TF in regulating ion channels expression in the heart

TBX5 drives expression of cardiac conduction system function

TBX5




rs 883079 eQTL



Arnolds et al. J Clin Invest. 2012

➤ Indirect regulation of Nav1.5 expression by TF


GWAS: identification of 3 loci associated with structural proteins





➤ new molecular mechanisms

MAPRE2 is a trafficking protein / regulator of microtubules

(microtubule plus-end binding protein (EB2))





eQTL



➤ EB1 protein, previously described in microtubules-mediated trafficking of connexin in cardiomyocytes

S. Xiao & R M Shaw, Trends Cardiovasc Med. 2015

MAPRE2 is a regulator of microtubule organization and impacts on sodium current



David Chiang
(Boston)




Mariam Jouni
(Chicago)




Modulation of microtubule function alters ion-channel trafficking

Multiple levels of sodium current dosage regulation affects in BrS phenotype




Normal sodium current

Rare coding mutations
(loss of function)




Reduced sodium current

Ion channel
control




Reduced sodium current

Translational
control



Reduced sodium current

Post-translational
control




Reduced sodium current

Common
non-coding
risk alleles


Impact in clinical practice: PRD and risk allele distribution among cases

Cumulative effect of susceptibility variants (21 loci)


Polygenic risk score



Polygenic Risk Score (PRS) in *SCN5A* +/-



BrS, a relevant model for more complex diseases



Genome-wide association meta-analysis identifies novel Brugada Syndrome susceptibility

loci and highlights multiple pathways modulating ion channel dosage



Julien Barc^{1*}, Rafik Tadros^{2,3*}, Charlotte Glinge^{2,4*}, David Y Chiang^{5*}, Mariam Jouni^{6*}, Floriane Simonet^{1*}, Sean J Jurgens⁷, Manon Baudic¹, Michele Nicastro², Franck Potet⁸, Joost A Offerhaus², Roddy Walsh², Seung Hoan Choi⁷, Arie O Verkerk^{2,9}, Yuka Mizusawa², Soraya Anys¹⁰, Damien Minois¹⁰, Marine Arnaud¹⁰, Josselin Duchateau^{11,12,13,14}, Yanushi D Wijeyeratne¹⁵, Alison Muir¹⁶, Michael Papadakis¹⁵, Silvia Castelletti¹⁷, Margherita Torchio¹⁸, Cristina Gil Ortuño¹⁹, Javier Lacunza²⁰, Daniela F Giachino^{21,22}, Natascia Cerrato²³, Raphaël P Martins²⁴, Oscar Campuzano^{25,26,27,28}, Sonia Van Dooren^{29,30}, Aurélie Thollet¹⁰, Florence Kyndt¹⁰, Andrea Mazzanti³¹, Nicolas Clémenty³², Arnaud Bisson³², Anniek Corveleyn³³, Birgit Stallmeyer³⁴, Sven Dittmann³⁴, Johan Saenen³⁵, Antoine Noël³⁶, Sherry Honarbakhsh³⁷, Boris Rudic^{38,39}, Halim Marzak⁴⁰, Matthew K Rowe⁴¹, Claire Federspiel⁴², Sophie Lepage⁴³, Leslie Placide⁴⁴, Antoine Milhem⁴⁵, Hector Barajas-Martinez⁴⁶, Britt-Maria Beckmann⁴⁷, Ingrid PC Krapels⁴⁸, Johannes Steinfurt⁴⁹, Bo Gregers Winkel⁵⁰, Reza Jabbari⁵¹, Moore B Shoemaker⁵², Carol Ann Remme², Bas J Boukens⁹, Doris Škorić-Milosavljević², Hennie Bikker⁵³, Federico C Manevy², Peter Lichtner⁵⁴, Marta Ribasés⁵⁵, Thomas Meitinger⁵⁴, Martina Müller-Nurasyid^{56,57,58,59}, KORA-Study Group⁶⁰, Jan H Veldink⁶¹, Leonard H van den Berg⁶¹, Philip Van Damme⁶², Daniele Cusi⁶³, Chiara Lanzani⁶⁴, Sidwell Rigade¹, Eric Charpentier^{1,65}, Estelle Baron¹, Stéphanie Bonnaud^{1,65}, Simon Lecoite¹, Audrey Donnart^{65,1}, Hervé Le Marec¹, Stéphanie Chatel¹, Matilde Karakachoff¹, Stéphane Bézieau¹⁰, Barry London⁶⁶, Jacob Tfelt-Hansen^{67,68,30}, Dan Roden^{69,70,71}, Katja E Odening^{49,72}, Marina Cerrone⁷³, Larry A Chinitz⁷³, Paul GA Volders⁷⁴, Maarten P van de Berg⁷⁵, Gabriel Laurent⁷⁶, Laurence Faivre⁷⁷, Charles Antzelevitch⁴⁶, Stefan Käb^{78,79}, Alain Al Arnaout⁴⁵, Jean-Marc Dupuis⁴³, Jean-Luc Pasquie⁸⁰, Olivier Billon⁴², Jason D Roberts⁴¹, Laurence Jesel^{40,81}, Martin Borggreffe^{38,39}, Pier D Lambiasi^{82,37}, Jacques Mansourati³⁶, Bart Loeys⁸³, Antoine Leenhardt^{84,30}, Pascale Guicheney^{85,86}, Philippe Maury⁸⁷, Eric Schulze-Bahr PhD^{34,30}, Tomas Robyns^{88,89}, Jeroen Breckpot³³, Dominique Babuty³², Silvia G Priori^{31,30}, Carlo Napolitano^{31,30}, Nantes Referral Center for inherited cardiac arrhythmia⁹⁰, Carlo de Asmundis^{91,92,27,30}, Pedro Brugada⁹³, Ramon Brugada⁹⁴, Elena Arbelo⁹⁵, Josep Brugada⁹⁶, Philippe Mabo²⁴, Nathalie Behar⁹⁷, Carla Giustetto²³, Maria Sabater Molina¹⁹, Juan R Gimeno^{20,30}, Can Hasdemir⁹⁸, Peter J Schwartz^{17,30}, Lia Crotti^{99,30}, Pascal P Mckeown¹⁰⁰, Sanjay Sharma¹⁵, Elijah R Behr^{15,30}, Michel Haissaguerre^{11,12,13,14}, Frédéric Sacher^{11,12,13,14}, Caroline Rooryck^{101,102}, Hanno L Tan¹⁰³, Pieter G Postema², Mario Delmar¹⁰⁴, Patrick T Ellinor¹⁰⁵, Steven A Lubitz¹⁰⁵, Jean-Baptiste Gourraud¹⁰, Michael WT Tanck¹⁰⁶, Alfred L George, Jr.¹⁰⁷, Calum A MacRae¹⁰⁸, Paul W BurrIDGE^{109,110}, Christian Dina¹, Vincent Probst^{10,30*}, Arthur A Wilde^{2,30*}, Jean-Jacques Schott^{10*}, Richard Redon^{10*}, Connie R Bezzina^{2*}

nature genetics

Barc et al.2022 Mar;54(3):232-239



Further perspectives (GWAS)




- 1- A first Genome-wide association meta-analysis **of Japanese ancestry**: 940 BrS cases and 1,634 controls
- 2- Genome-wide association meta-analysis combining the Japanese and the European-ancestry datasets: **3,760 BrS cases** and 11,635 controls
 - **17 loci** reached the genome-wide significance
 - **5 loci** were newly discovered in the cross-ancestry meta-analysis.
 - **Polygenic risk score analysis highlights shared genetic architecture across ancestries**
- 3- **Genome-wide analysis of lethal arrhythmic events among BrS patients identifies 1 VF-associated SNP in Japanese and European BrS**

Whole genome sequencing accurately identifies CNVs



→ ECG de type I Brugada


3 syncopes → ICD at age 45




50% reduction of coverage
→ hétérozygote deletion

➤ 10 kb deletion in the promoter region of *SCN5A*


Functional annotation




Functional annotation of the deletion region



Functional annotation of the deletion region



Functional annotation of the 10kb deletion (patch clamping)



➤ Reduction of I_{Na} density

Key take-home messages

- The inheritance model for Brugada syndrome is complex
- Unexpected large effect of common genetic variations on BrS susceptibility
- **The *SCN5A* locus is prominent in disease susceptibility, with strong involvement in both rare and common alleles**
- Strong involvement of transcription regulation impacting ion channels expression, heart development **as well as cardiac structural anomalies**